

2. **H. non-scripta** (L.) Chouard ex Rothm. (Endymion non-scriptus (L.) Garcke) - Bluebell. Stems to 50cm; leaves up to 20mm wide; racemes pendent at apex, 1-sided, with pendent strongly sweetly scented flowers; tepals 14-20mm, forming \pm parallel-sided tubular perianth, strongly recurved at apex, outer 3 stamens fused to perianth for $>3/4$ their length. Native; woods, hedgerows, shady banks, grassland in wetter regions; frequent to abundant throughout BI.

2 x 3. **H. non-scripta** x **H. hispanica** is more commonly grown in gardens and natd than H. hispanica and arises naturally where natd H. hispanica meets native or natd H. non-scripta; frequent in Br and CI in similar places to H. hispanica. It is intermediate in all characters and fertile, forming a complete spectrum between the parents and often natd in absence of both.

3. **H. hispanica** (Miller) Rothm. (Endymion hispanicus (Miller) Chouard) - Spanish Bluebell. Stems to 40cm; leaves up to 35mm wide; racemes erect, not 1-sided, with erect to patent, faintly scented flowers; tepals 12-18mm, forming campanulate perianth, not recurved at apex; outer 3 stamens fused to perianth for $<3/4$ their length. Intrd; grown in gardens, natd in woods, copses, shady banks and field-borders; scattered in Br and CI; Spain and Portugal. Over-recorded for H. hispanica x H. non-scripta.